

August 30, 2012

Recent results for D^0 - $\bar{D}^0$ mixing and CP violation, and HFAG averages

NICOLA NERI¹

*Istituto Nazionale di Fisica Nucleare
Sezione di Milano, I-20131 Milano, ITALY*

I present here the HFAG averages for the parameters that regulate flavor mixing and CP violation in the neutral D meson system. I also discuss recent results from the B factories for the measurements of the mixing parameter y_{CP} and the CP violation parameter ΔY (A_Γ) in the lifetime ratio analysis of the transitions to the CP -even eigenstates $D^0 \rightarrow K^+K^-, \pi^+\pi^-$, relative to the transitions to the CP -mixed state $D^0 \rightarrow K^-\pi^+$.

PRESENTED AT

The 5th International Workshop on Charm Physics
(Charm 2012)
Honolulu, Hawai'i (USA), May 14-17, 2012

¹on behalf of the *BABAR* collaboration.

1 Introduction

Several measurements have reported evidence for flavor mixing in the neutral D meson system [1, 2, 3, 4, 5, 6]. The results are in agreement with the Standard Model (SM) predictions, which unfortunately are affected by large theoretical uncertainties [7, 8, 9]. However, the increasing precision of D^0 - $\bar{D}^0$ mixing measurements helps constrain new physics models [10, 11]. CP violation in D meson decays, though notoriously difficult to calculate precisely, is expected to be very small in the SM, at the level of 10^{-3} or less [8, 12, 13]. Relatively large CP asymmetries, at the percent level, might be a signature of new physics effects. Recent results from the LHCb experiment [14] reported evidence for direct CP violation measuring the difference of CP asymmetries in singly-Cabibbo suppressed $D^0 \rightarrow \pi^+\pi^-$ * and $D^0 \rightarrow K^+K^-$ decays, with a statistical significance of 3.5σ . The observed asymmetries are marginally compatible with the SM but not conclusive for establishing new physics [15, 16, 17]. These intriguing results renew interest in studying mixing and CP violation in the $D^0\bar{D}^0$ meson system and in general in the charm physics sector.

The Heavy Flavor Averaging Group (HFAG) [18] provides averages for heavy flavor quantities. It is divided into several sub-groups, each of which focuses on a different set of heavy flavor measurements. The Charm Physics sub-group [19] studies the following topics: D^0 - $\bar{D}^0$ mixing, CP violation, spectroscopy of charm mesons and baryons, semileptonic decays, decay constants, hadronic branching fraction measurements and rare D decay modes. In particular, it provides averages for the the D^0 - $\bar{D}^0$ mixing and CPV parameters by combining measurements from different experiments in a χ^2 -based fit. The results of the fit are expressed in term of physics parameters that can be directly compared to the theoretical predictions.

2 HFAG notations for flavor mixing and CPV

Flavor mixing occurs when the Hamiltonian eigenstates (or physics eigenstates) D_1 , D_2 differ from the flavor eigenstates D^0 , $\bar{D}^0$. A neutral D meson produced at time $t = 0$ in a definite flavor state D^0 , will then evolve and oscillate into a state of opposite flavor $\bar{D}^0$, after a certain time. If we describe the time evolution of the flavor eigenstates in terms of a 2×2 effective Hamiltonian, $H = M - \frac{i}{2}\Gamma$, then, assuming CPT is conserved, the mass eigenstates can be expressed in terms of the flavor eigenstates by,

$$\begin{aligned} |D_1\rangle &= p|D^0\rangle + q|\bar{D}^0\rangle \\ |D_2\rangle &= p|D^0\rangle - q|\bar{D}^0\rangle \end{aligned} \tag{1}$$

*The use of charge conjugate reactions is implied throughout unless stated otherwise.

with the normalization $|q|^2 + |p|^2 = 1$ and

$$\frac{q}{p} = \sqrt{\frac{M_{12}^* - i/2\Gamma_{12}^*}{M_{12} - i/2\Gamma_{12}}}. \quad (2)$$

Assuming a phase convention such that $CP|D^0\rangle = -|\bar{D}^0\rangle$ and $CP|\bar{D}^0\rangle = -|D^0\rangle$ then, if CP is conserved, we have that $q = p = 1/\sqrt{2}$ and the mass eigenstates coincide with the CP eigenstates: $|D_1\rangle = |D_{CP-}\rangle$ (CP -odd) and $|D_2\rangle = |D_{CP+}\rangle$ (CP -even).

The mixing parameters can be expressed in terms of the difference of masses ($m_{1,2}$) and widths ($\Gamma_{1,2}$) of the Hamiltonian eigenstates,

$$x = \frac{m_2 - m_1}{\Gamma}, \quad y = \frac{\Gamma_2 - \Gamma_1}{2\Gamma}, \quad (3)$$

where $\Gamma = (\Gamma_1 + \Gamma_2)/2$.

CP violation can be of three types:

1. *CPV in decay or direct CPV*: this occurs when the decay amplitudes for CP conjugate processes are different in modulus. If $\langle f|H|D^0\rangle = A_f$, $\langle \bar{f}|H|\bar{D}^0\rangle = \bar{A}_{\bar{f}}$ are the D^0 and $\bar{D}^0$ decay amplitudes into the final states f and CP conjugate $\bar{f}$, then

$$\left| \frac{\bar{A}_{\bar{f}}}{A_f} \right| \neq 1 \implies CPV. \quad (4)$$

2. *CPV in mixing or indirect CPV*: it occurs when the Hamiltonian eigenstates do not coincide with the CP eigenstates. That is

$$\left| \frac{q}{p} \right| \neq 1 \implies CPV. \quad (5)$$

3. *CPV in the interference of mixing and decay*: for neutral D mesons there is a third possibility to observe CP violation even when CP is conserved in mixing and also in decay. In this case, CP violation arises when, in a process with final state f that can be reached by neutral D mesons of both flavors (*i.e.* D^0 and $\bar{D}^0$), there is a relative weak phase[†] difference between the mixing and the decay amplitudes. The quantity of interest that is independent of phase conventions, and physically meaningful, is

$$\lambda_f \equiv \frac{q}{p} \frac{\bar{A}_f}{A_f} \equiv \left| \frac{q}{p} \right| \left| \frac{\bar{A}_f}{A_f} \right| e^{i(\delta_f + \phi_f)}, \quad (6)$$

[†]The CP -violating phase is also indicated as a weak phase since it originates from the weak interaction in the SM.

where δ_f and ϕ_f are the CP -conserving and CP -violating phases respectively. If CP is conserved in mixing and in decay, the signature of CP violation in the interference of mixing and decay is thus

$$\phi_f \neq 0 \implies CPV. \quad (7)$$

For CP eigenstates, CPV in either mixing or decay is indicated by

$$|\lambda_f| \neq 1, \quad (8)$$

while CPV in the interference of mixing and decay corresponds to

$$\text{Im}(\lambda_f) \neq 1. \quad (9)$$

Note that if there is no weak phase in the decay amplitudes then $\arg(q/p) = \phi$ and it is independent of the final state f .

3 Experimental observables and parameters of the theory

The most precise constraints on the mixing parameters x , y and the CPV parameters $|q/p|$, ϕ are obtained in time-dependent analyses of D decays.

Consider a final state f that can be reached by both D^0 and $\bar{D}^0$ decays. If a D^0 is produced at time $t = 0$, it can reach the final state f by mixing to $\bar{D}^0$ followed by the decay $\bar{D}^0 \rightarrow f$, or directly through the decay $D^0 \rightarrow f$. The interference between the mixing and decay amplitudes modifies the time-dependence with respect to the pure exponential as follows:

$$\begin{aligned} \frac{|\langle f | H | D^0(t) \rangle|^2}{\frac{1}{2}e^{-\Gamma t}} &= \left(|A_f|^2 + \left| \frac{q}{p} \right|^2 |\bar{A}_f|^2 \right) \cosh(y\Gamma t) + 2\text{Re} \left(\frac{q}{p} A_f^* \bar{A}_f \right) \sinh(y\Gamma t) \\ &+ \left(|A_f|^2 - \left| \frac{q}{p} \right|^2 |\bar{A}_f|^2 \right) \cos(x\Gamma t) - 2\text{Im} \left(\frac{q}{p} A_f^* \bar{A}_f \right) \sin(x\Gamma t) \quad (10) \end{aligned}$$

$$\begin{aligned} \frac{|\langle f | H | \bar{D}^0(t) \rangle|^2}{\frac{1}{2}e^{-\Gamma t}} &= \left(|\bar{A}_f|^2 + \left| \frac{p}{q} \right|^2 |A_f|^2 \right) \cosh(y\Gamma t) + 2\text{Re} \left(\frac{p}{q} A_f \bar{A}_f^* \right) \sinh(y\Gamma t) \\ &+ \left(|\bar{A}_f|^2 - \left| \frac{p}{q} \right|^2 |A_f|^2 \right) \cos(x\Gamma t) - 2\text{Im} \left(\frac{p}{q} A_f \bar{A}_f^* \right) \sin(x\Gamma t). \quad (11) \end{aligned}$$

HFAG combines 38 observables measured in time-dependent and time-integrated analyses from the following experiments: *BABAR*, Belle, CDF, CLEO, CLEOc, E791, FOCUS, and LHCb. When allowing for CPV there are 10 underlying parameters that are

extracted from the χ^2 fit: $x, y, |q/p|, \phi, \delta, \delta_{K\pi\pi}, R_D, A_D, A_\pi, A_K$. The parameters $\delta, \delta_{K\pi\pi}$ are relative strong phases, R_D is the ratio $\Gamma(D^0 \rightarrow K^+\pi^-)/\Gamma(\bar{D}^0 \rightarrow K^-\pi^+)$ and A_D, A_π, A_K are the direct CP -violation asymmetries for the $D^0 \rightarrow K^+\pi^-$, $D^0 \rightarrow \pi^+\pi^-$ and $D^0 \rightarrow K^+K^-$ modes. The relationships between these parameters and the measured observables are given below.

1. *Semileptonic decays*: search for mixing by reconstructing the “wrong-sign” (WS) decay chain, $D^{*+} \rightarrow D^0\pi^+, D^0 \rightarrow \bar{D}^0, \bar{D}^0 \rightarrow K^{(*)+}e^-\bar{\nu}_e$. In contrast to hadronic decays, the WS charge combinations can occur only through mixing. The measurement of R_M is related to the mixing parameters as follows

$$R_M = \frac{1}{2}(x^2 + y^2), \quad (12)$$

and can be obtained directly as the ratio of WS to right-sign (RS) signal events. The RS events correspond to the non-mixed process.

2. *Decays to CP eigenstates*: measure the mixing parameter y_{CP} and the CPV parameter A_Γ with a lifetime ratio analysis of the transitions to the CP eigenstates and the transitions to the CP -mixed state $D^0 \rightarrow K^-\pi^+$,

$$\begin{aligned} 2y_{CP} &= \left(\left| \frac{q}{p} \right| + \left| \frac{p}{q} \right| \right) y \cos \phi - \left(\left| \frac{q}{p} \right| - \left| \frac{p}{q} \right| \right) x \sin \phi \\ 2A_\Gamma &= \left(\left| \frac{q}{p} \right| - \left| \frac{p}{q} \right| \right) y \cos \phi - \left(\left| \frac{q}{p} \right| + \left| \frac{p}{q} \right| \right) x \sin \phi. \end{aligned} \quad (13)$$

The parameter A_Γ is the decay-rate asymmetry for the CP eigenstates. If CP is conserved $y_{CP} = y$ and $A_\Gamma = 0$.

Time-integrated CP -violating asymmetries in CP -even eigenstates, *e.g.* in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$, provide constraints on the mixing and CPV parameters according to the relations,

$$\begin{aligned} \frac{\Gamma(D^0 \rightarrow K^+K^-) + \Gamma(\bar{D}^0 \rightarrow K^-K^+)}{\Gamma(D^0 \rightarrow K^-K^+) + \Gamma(\bar{D}^0 \rightarrow K^+K^-)} &= A_K + \frac{\langle t \rangle}{\tau_D} A_{CP}^{\text{indirect}} \\ \frac{\Gamma(D^0 \rightarrow \pi^+\pi^-) - \Gamma(\bar{D}^0 \rightarrow \pi^-\pi^+)}{\Gamma(D^0 \rightarrow \pi^-\pi^+) + \Gamma(\bar{D}^0 \rightarrow \pi^+\pi^-)} &= A_\pi + \frac{\langle t \rangle}{\tau_D} A_{CP}^{\text{indirect}}, \end{aligned} \quad (14)$$

where

$$2A_{CP}^{\text{indirect}} = \left(\left| \frac{q}{p} \right| + \left| \frac{p}{q} \right| \right) x \sin \phi - \left(\left| \frac{q}{p} \right| - \left| \frac{p}{q} \right| \right) y \cos \phi, \quad (15)$$

$\langle t \rangle$ is the average reconstructed D^0 proper time and τ_D is the nominal D^0 lifetime.

3. *Three-body $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $D^0 \rightarrow K_S^0 K^+ K^-$ decays:* measure directly the mixing and *CPV* parameters x , y , $|q/p|$, and ϕ with a time-dependent Dalitz plot analysis.
4. *Wrong-sign decays to hadronic non-CP eigenstates:* measure the parameters $x'^{\pm}$ and $y'^{\pm}$ and R_D and A_D in a time-dependent analysis of the WS events selected through the decay chain $D^{*+} \rightarrow D^0 \pi^+$, $D^0 \rightarrow K^+ \pi^-$. The parameters are defined as follows:

$$\begin{aligned}
x'^{\pm} &= \left(\frac{1 \pm A_M}{1 \mp A_M} \right)^{1/4} (x' \cos \phi \pm y' \sin \phi) \\
y'^{\pm} &= \left(\frac{1 \pm A_M}{1 \mp A_M} \right)^{1/4} (y' \cos \phi \mp x' \sin \phi) \\
\frac{\Gamma(D^0 \rightarrow K^+ \pi^-) + \Gamma(\bar{D}^0 \rightarrow K^- \pi^+)}{\Gamma(D^0 \rightarrow K^- \pi^+) + \Gamma(\bar{D}^0 \rightarrow K^+ \pi^-)} &= R_D \\
\frac{\Gamma(D^0 \rightarrow K^+ \pi^-) - \Gamma(\bar{D}^0 \rightarrow K^- \pi^+)}{\Gamma(D^0 \rightarrow K^- \pi^+) + \Gamma(\bar{D}^0 \rightarrow K^+ \pi^-)} &= A_D
\end{aligned} \tag{16}$$

where

$$\begin{aligned}
\begin{pmatrix} x' \\ y' \end{pmatrix} &= \begin{pmatrix} \cos \delta & \sin \delta \\ -\sin \delta & \cos \delta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\
A_M = \frac{|q/p|^2 - |p/q|^2}{|q/p|^2 + |p/q|^2}, & \quad \delta = \arg \left(\frac{A(D^0 \rightarrow K^+ \pi^-)}{A(\bar{D}^0 \rightarrow K^+ \pi^-)} \right).
\end{aligned}$$

The superscript in $x'^{\pm}$ and $y'^{\pm}$ identifies the flavor of the D sample, *i.e.* D^0 (+) and $\bar{D}^0$ (-).

The mixing parameters x'' and y'' ,

$$\begin{pmatrix} x'' \\ y'' \end{pmatrix} = \begin{pmatrix} \cos \delta_{K\pi\pi} & \sin \delta_{K\pi\pi} \\ -\sin \delta_{K\pi\pi} & \cos \delta_{K\pi\pi} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

may be measured by means of a time-dependent Dalitz plot analysis of the three-body WS events $D^0 \rightarrow K^+ \pi^- \pi^0$. The strong phase that rotates the mixing parameters x and y is defined here as $\delta_{K\pi\pi} = \arg \left(\frac{A(D^0 \rightarrow K^+ \pi^- \pi^0)}{A(\bar{D}^0 \rightarrow K^+ \pi^- \pi^0)} \right)$.

5. $\psi(3770) \rightarrow D^0 \bar{D}^0$ reaction: exploiting the quantum correlations of the $D^0 \bar{D}^0$ production at threshold, it is possible to constrain the physics parameters by measuring x^2 , y , R_D , $2\sqrt{R_D} \cos \delta$, $2\sqrt{R_D} \sin \delta$.

4 HFAG results for D^0 - $\bar{D}^0$ mixing and CPV

HFAG uses a global fit to determine world averages of mixing parameters, CPV parameters, and strong phases. The fit uses 38 observables taken from measurements of $D^0 \rightarrow K^+\ell^-\nu$, $D^0 \rightarrow K^+K^-$, $D^0 \rightarrow \pi^+\pi^-$, $D^0 \rightarrow K^+\pi^-$, $D^0 \rightarrow K^+\pi^-\pi^0$, $D^0 \rightarrow K_S^0\pi^+\pi^-$, $D^0 \rightarrow K_S^0K^+K^-$ decays and from double-tagged branching fractions measured at the $\psi(3770)$ resonance. Correlations among observables are accounted for by using covariance matrices provided by the experimental collaborations. Errors are assumed to be Gaussian, and systematic errors among different experiments are assumed uncorrelated unless specific correlations have been identified. An independent log-likelihood fit which accounts for non-Gaussian errors has been used as a control check, and equivalent results have been obtained.

Averages for the parameters R_M , y_{CP} and A_Γ are calculated and then provided as input to the global fit. The observable R_M is calculated from $D^0 \rightarrow K^+\ell^-\nu$ decays [20, 21, 22, 23], and the average value is $R_M = (0.013 \pm 0.027)\%$. The inputs used for this average are plotted in Fig. 1. The observable y_{CP} is calculated from $D^0 \rightarrow K^+K^-/\pi^+\pi^-$ and from $D^0 \rightarrow K_S^0K^+K^-$ decays while A_Γ is calculated from $D^0 \rightarrow K^+K^-/\pi^+\pi^-$ decays only. The inputs used for these averages are plotted in Fig. 2. The average values are $y_{CP} = (1.064 \pm 0.209)\%$ and $A_\Gamma = (0.026 \pm 0.231)\%$.

The $D^0 \rightarrow K^+\pi^-$ observables used are from Belle [24], BABAR [1], and CDF [6]; earlier measurements have much less precision and are not used. The observables from $D^0 \rightarrow K_S^0\pi^+\pi^-$ decays for no- CPV are from Belle [25] and BABAR [26], but for the CPV -allowed case only Belle measurements [25] are available. The $D^0 \rightarrow K^+\pi^-\pi^0$ results are from BABAR [3], and the $\psi(3770) \rightarrow D^0\bar{D}^0$ results are from CLEOc [27]. Time-integrated measurements of CP -violating asymmetries (A_{CP}) in the $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ modes are from BABAR [29] and Belle [30]. The difference of the CP -violating asymmetries defined as $\Delta A_{CP} = A_{CP}(K^+K^-) - A_{CP}(\pi^+\pi^-)$ are from LHCb [14] and CDF [31]. These measurements reported evidence for direct CPV with a statistical significance of 3.5σ and 2.7σ , respectively.

Three types of fit are produced by HFAG with different CPV assumptions:

1. *No CPV* : in this fit it is assumed that CP is conserved and the CPV parameters A_D , A_K , A_π , $|q/p|-1$ and ϕ are fixed to zero.
2. *No direct CPV* : the CPV parameters A_D , A_K , A_π are fixed to zero. In this case the relation [32, 33] $\tan \phi = (1 - |q/p|^2)/(1 + |q/p|^2) \times (x/y)$ is satisfied, and this reduces the four independent parameters $(x, y, |q/p|, \phi)$ to three. The independent parameters used in the fit are $x_{12} = 2|M_{12}|/\Gamma$, $y_{12} = \Gamma_{12}/\Gamma$, and $\phi_{12} = \text{Arg}(M_{12}/\Gamma_{12})$, where M_{12} and Γ_{12} are the off-diagonal elements of the $D^0 - \bar{D}^0$ mass and decay matrices, respectively.




Figure 1: World average value of R_M from Ref. [19], as calculated from $D^0 \rightarrow K^+ \ell^- \nu$ measurements [20, 21, 22, 23].

3. *CPV-allowed*: where all the parameters for mixing and *CPV* are floated in the fit.

All fit results are listed in Table 1. The total χ^2 is 35.6 for $37 - 10 = 27$ degrees of freedom; this corresponds to a confidence level of 12.4%. The resulting 1σ - 5σ contours are shown in Fig. 3 for the *CP*-conserving case, in Fig. 4 for the no-direct-*CPV* case, and in Fig. 5 for the *CPV*-allowed case. For the *CPV*-allowed fit, the no-mixing point $(x, y) = (0, 0)$ is excluded at a confidence level of 1.28×10^{-24} corresponding to a statistical significance of 10.2σ . The parameter x differs from zero by 2.7σ , and y differs from zero by 6.0σ . In the $(|q/p|, \phi)$ plot, the point $(1, 0)$ is within the 1σ contour; thus the data are consistent with *CP* conservation in mixing and in the interference between mixing and decay.




Figure 2: World average value of y_{CP} and A_F from Ref. [19], as calculated from $D^0 \rightarrow K^+ K^- / \pi^+ \pi^-$.




Figure 3: Two-dimensional contours for the mixing parameters (x, y) , for no CPV .




Figure 4: Two-dimensional contours for theory parameters (x_{12}, y_{12}) (top left), (x_{12}, ϕ_{12}) (top right), and (y_{12}, ϕ_{12}) (bottom), for no direct *CPV*.




Figure 5: Two-dimensional contours for the parameters (x, y) (left) and $(|q/p|, \phi)$ (right), allowing for CPV .

Table 1: Results of the global fit for different assumptions concerning CPV .

Parameter	No CPV	No direct CPV	CPV -allowed	CPV -allowed 95% C.L.
x (%)	$0.65^{+0.18}_{-0.19}$	0.62 ± 0.19	$0.63^{+0.19}_{-0.20}$	[0.24, 0.99]
y (%)	0.73 ± 0.12	0.75 ± 0.12	0.75 ± 0.12	[0.51, 0.98]
δ ($^\circ$)	$21.0^{+9.8}_{-11.0}$	$22.2^{+9.9}_{-11.2}$	$22.1^{+9.7}_{-11.1}$	[-2.6, 40.6]
R_D (%)	0.3307 ± 0.0080	0.3305 ± 0.0080	0.3311 ± 0.0081	[0.315, 0.347]
A_D (%)	0.0	0.0	-1.7 ± 2.4	[-6.4, 3.0]
$ q/p $	1.0	$1.04^{+0.07}_{-0.06}$	$0.88^{+0.18}_{-0.16}$	[0.59, 1.26]
ϕ ($^\circ$)	0.0	$-2.02^{+2.67}_{-2.74}$	$-10.1^{+9.5}_{-8.9}$	[-27.4, 8.7]
$\delta_{K\pi\pi}$ ($^\circ$)	$17.8^{+21.7}_{-22.8}$	$19.4^{+21.8}_{-22.9}$	$19.3^{+21.8}_{-22.9}$	[-26.3, 61.8]
A_π	0.0	0.0	0.36 ± 0.25	[-0.13, 0.86]
A_K	0.0	0.0	-0.31 ± 0.24	[-0.78, 0.15]
x_{12} (%)	—	0.62 ± 0.19	—	—
y_{12} (%)	—	0.75 ± 0.12	—	—
ϕ_{12} ($^\circ$)	—	$4.9^{+7.7}_{-6.5}$	—	—

5 New results from the B factories

5.1 Measurement of D^0 - $\bar{D}^0$ mixing and search for indirect CP violation in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays

The *BABAR* [34] and the Belle experiments [35] have recently presented updated results for the measurements of the mixing parameter y_{CP} and the CP violation parameters ΔY [‡] for *BABAR* and A_Γ for Belle. The definitions of ΔY and A_Γ are the following:

$$\Delta Y = \frac{\Gamma^+ - \bar{\Gamma}^+}{2\Gamma}, \quad A_\Gamma = \frac{\bar{\tau}^+ - \tau^+}{\bar{\tau}^+ + \tau^+} \quad \text{and} \quad \Delta Y = (1 + y_{CP})A_\Gamma, \quad (17)$$

where $\tau^+ = 1/\Gamma^+$ ($\bar{\tau}^+ = 1/\bar{\Gamma}^+$) are the effective lifetimes for D^0 ($\bar{D}^0$) decaying to the CP -even final states K^+K^- and $\pi^+\pi^-$. In principle the parameters y_{CP} and ΔY depend on the final state f , as indicated in Eq. (18) and Eq. (19) below, when accounting for direct CP violation. The *BABAR* and Belle analyses assume CP conservation in the decay (*i.e.* $A_\pi = 0$, $A_K = 0$), and neglect terms of $\mathcal{O}(10^{-4})$ in the expressions below, that are beyond the present experimental sensitivity:

$$y_{CP}^f = y \cos \phi_f + \frac{1}{2}(A_M + A_f)x \sin \phi_f - \frac{1}{4}A_M A_f y \cos \phi_f \quad (18)$$

$$\Rightarrow y_{CP} \simeq y \cos \phi + \frac{1}{2}A_M x \sin \phi$$

$$\Delta Y^f = -x \sin \phi_f + \frac{1}{2}(A_M + A_f)y \cos \phi_f + \frac{1}{4}A_M A_f x \sin \phi_f \quad (19)$$

$$\Rightarrow \Delta Y \simeq -x \sin \phi + \frac{1}{2}A_M y \cos \phi.$$

Hence, the parameters no longer depend on the final state f , and so can be averaged over the K^+K^- and $\pi^+\pi^-$ modes.

The measurements are based on the ratio of lifetimes extracted simultaneously from a sample of D^0 mesons produced through the flavor-tagged process $D^{*+} \rightarrow D^0\pi^+$, where the D^0 decays to $K^-\pi^+$, K^-K^+ , or $\pi^-\pi^+$; *BABAR* uses the additional samples of untagged decays $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^-K^+$ for the measurement of y_{CP} . These have about 4 times the statistics of the corresponding flavor-tagged sample, but have lower purity.

The main selection criteria require that the center-of-mass momentum of the D^0 (p^*) be greater than 2.5 GeV/ c [§], that D^0 daughter tracks be identified as kaons and pions, and require for the flavor-tagged sample that the reconstructed $\Delta m =$

[‡]Note that this definition for ΔY has different sign convention with respect to *BABAR* results published previously [2, 4].

[§]Belle requires $p^* > 3.1$ GeV/ c for the data collected at the $\Upsilon(5S)$ mass peak.

Category	Δy_{CP} (%)	$\Delta(\Delta Y)$ (%)
Fit region	0.057	0.022
Signal model	0.022	0.000
Charm bkg	0.045	0.001
Combinatorial bkg	0.079	0.002
Selection	0.059	0.054
Total	0.124	0.058

Table 2: Systematic uncertainties for y_{CP} and ΔY at *BABAR*. The total is the sum-in-quadrature of the entries in each column.

$m(D^{*+}) - m(D^0)$ be close to the value of 0.1455 GeV. The selection criterion on p^* is used to reject D mesons from B decays, and to improve the signal significance.

The proper time t and proper-time error σ_t are obtained from the reconstruction of the 3-dimensional flight length ($\vec{L}$) and the momentum of the D^0 ($\vec{p}$) according to the relation $t = m/|\vec{p}|^2 \vec{L} \cdot \vec{p}$, where m is the nominal mass of the D^0 . The flight length is reconstructed by means of a kinematic fit to the decay vertex and production vertex of the D^0 , the latter being constrained to originate within the e^+e^- collision region. The typical transverse dimensions of the luminous region of the PEP-II collider are about 100 μm in the x direction and 7 μm in the y direction. The most probable σ_t value is about 40% of the nominal D^0 lifetime, and only candidates with $\sigma_t < 0.5$ ps are retained for the fit in the *BABAR* analysis.

The lifetimes of the CP -even modes K^-K^+ , $\pi^-\pi^+$ are compared to that of the CP -mixed mode $K^-\pi^+$ in order to measure y_{CP} , which is proportional to the ratio of the lifetimes, and ΔY (A_Γ) which is proportional to the difference of the effective lifetimes of $\bar{D}^0$ and D^0 into CP -even modes. *BABAR* measures $y_{CP} = [0.72 \pm 0.18(\text{stat}) \pm 0.12(\text{syst})]\%$ and $\Delta Y = [0.09 \pm 0.26(\text{stat}) \pm 0.06(\text{syst})]\%$ using a data sample corresponding to an integrated luminosity of 468 fb^{-1} [34]. Belle measures $y_{CP} = [1.11 \pm 0.22(\text{stat}) \pm 0.11(\text{syst})]\%$ and $A_\Gamma = [-0.03 \pm 0.20(\text{stat}) \pm 0.08(\text{syst})]\%$ using a data sample corresponding to 976 fb^{-1} [35].

The systematic uncertainties on y_{CP} and ΔY are reported in Table 2 for *BABAR*, and on y_{CP} and A_Γ are reported in Table 3 for Belle. The total systematic uncertainties are comparable between the two experiments. It is worth noting that the Belle experiment quotes a systematic error due to the silicon vertex detector (SVD) misalignment that is negligible in the case of the *BABAR* experiment. The measurement of the proper time average value in Belle shows a dependence on the cosine of the polar angle in the e^+e^- center-of-mass ($\cos\theta^*$) that is not properly reproduced by Monte Carlo simulation. The discrepancy between data and Monte Carlo is different for the data collected with the two vertex detectors that have been used during the running of the experiment. In order to minimize the systematic error due to this

Category	Δy_{CP} (%)	ΔA_Γ (%)
Acceptance	0.050	0.044
SVD misalignments	0.060	0.041
Mass window position	0.007	0.009
Background	0.059	0.050
Resolution function	0.030	0.002
Total	0.11	0.08

Table 3: Systematic uncertainties for y_{CP} and A_Γ at Belle. The total is the sum-in-quadrature of the entries in each column.

effect, Belle has performed the measurement by dividing the sample into different intervals of $\cos \theta^*$.

The y_{CP} measurements report evidence for D^0 - $\bar{D}^0$ mixing with a significance of 3.3σ in the case of *BABAR* (most precise measurement to date) and 4.5σ in the case of Belle. The measurements of ΔY from *BABAR*, and A_Γ from Belle are compatible within error and both are consistent with no *CPV*. The new measurements supersede the previous *BABAR* [2, 4] and Belle results [5]. The updated HFAG averages, including the new results, are $y_{CP} = (0.866 \pm 0.155)\%$ and $A_\Gamma = (-0.022 \pm 0.161)\%$. The comparison with the previous HFAG average values $y_{CP} = (1.064 \pm 0.209)\%$ and $A_\Gamma = (0.026 \pm 0.231)\%$ indicates significant improvement in precision and a lower central value for y_{CP} .

6 Summary

In summary, HFAG provides averages for the D^0 - $\bar{D}^0$ mixing and *CP*-violating parameters by combining results from the following experiments: *BABAR*, Belle, CDF, CLEO, CLEOc, E791, FOCUS, and LHCb. The average values from the *CPV*-allowed fit for the mixing parameters are $x = (0.63^{+0.19}_{-0.20})\%$, $y = (0.72 \pm 0.12)\%$, and for the *CP*-violating parameters are $|q/p| = 0.88^{+0.18}_{-0.16}$, $\phi = (-10.1^{+9.5}_{-8.9})^\circ$. These average values do not include the new *BABAR* and Belle results presented here. The parameter x differs from zero by 2.7σ , and y differs from zero by 6.0σ . The no-mixing hypothesis, ($x = 0, y = 0$), is excluded with a statistical significance of 10.2σ . The LHCb and CDF experiments have obtained first evidence for direct *CPV* in D^0 decays, while there is no evidence for *CPV* in mixing ($|q/p| \neq 1$) and in the interference between mixing and decay ($\phi \neq 0$).

New results for D^0 - $\bar{D}^0$ mixing and *CPV* in the lifetime ratio analysis of the transitions $D^0 \rightarrow K^+K^-/\pi^+\pi^-$ to the transitions $D^0 \rightarrow K^-\pi^+$ have been presented. *BABAR* measures $y_{CP} = [0.72 \pm 0.18(\text{stat}) \pm 0.12(\text{syst})]\%$ and $\Delta Y = [0.09 \pm$

$0.26(\text{stat}) \pm 0.06(\text{syst})\%$ using a data sample corresponding to an integrated luminosity of 468 fb^{-1} . Belle measures $y_{CP} = [1.11 \pm 0.22(\text{stat}) \pm 0.11(\text{syst})]\%$ and $A_\Gamma = [-0.03 \pm 0.20(\text{stat}) \pm 0.08(\text{syst})]\%$ using a data sample corresponding to 976 fb^{-1} . The updated HFAG averages, including the new results, are $y_{CP} = (0.866 \pm 0.155)\%$ and $A_\Gamma = (-0.022 \pm 0.161)\%$.

HFAG averages for D^0 - $\bar{D}^0$ mixing and CPV parameters are in agreement with the SM. Recent results for direct CPV in $D^0 \rightarrow K^+ K^- / \pi^+ \pi^-$ decays report a larger asymmetry value than the SM expectation. However, given the present knowledge of the charm system, this cannot (yet) be considered a clear signal of physics beyond the SM.

ACKNOWLEDGEMENTS

I am grateful to my *BABAR* colleagues for providing new results in time for the workshop and for useful comments. I would like to thank the conference organizers for their warm hospitality in a wonderful location.

References

- [1] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **98**, 211802 (2007).
- [2] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **78**, 011105 (2008).
- [3] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **103**, 211801 (2009).
- [4] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **80**, 071103 (2009).
- [5] M. Staric *et al.* [Belle Collaboration], Phys. Rev. Lett. **98**, 211803 (2007).
- [6] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **100**, 121802 (2008).
- [7] Z. -z. Xing, Phys. Rev. D **55**, 196 (1997).
- [8] S. Bianco, F. L. Fabbri, D. Benson and I. Bigi, Riv. Nuovo Cim. **26N7**, 1 (2003).
- [9] G. Burdman and I. Shipsey, Ann. Rev. Nucl. Part. Sci. **53**, 431 (2003).
- [10] E. Golowich, J. Hewett, S. Pakvasa and A. A. Petrov, Phys. Rev. D **76**, 095009 (2007).
- [11] E. Golowich, J. Hewett, S. Pakvasa and A. A. Petrov, Phys. Rev. D **79**, 114030 (2009).

- [12] F. Buccella, M. Lusignoli, G. Miele, A. Pugliese and P. Santorelli, Phys. Rev. D **51**, 3478 (1995).
- [13] Y. Grossman, A. L. Kagan and Y. Nir, Phys. Rev. D **75**, 036008 (2007).
- [14] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **108**, 111602 (2012).
- [15] Y. Grossman, A. L. Kagan and J. Zupan, arXiv:1204.3557 [hep-ph].
- [16] E. Franco, S. Mishima and L. Silvestrini, JHEP **1205**, 140 (2012).
- [17] G. Isidori, J. F. Kamenik, Z. Ligeti and G. Perez, Phys. Lett. B **711**, 46 (2012).
- [18] HFAG web page: <http://www.slac.stanford.edu/xorg/hfag/>
- [19] HFAG, Charm Physics sub-group: <http://www.slac.stanford.edu/xorg/hfag/charm/index.html>
- [20] E. M. Aitala *et al.* [E791 Collaboration], Phys. Rev. Lett. **77**, 2384 (1996).
- [21] C. Cawfield *et al.* [CLEO Collaboration], Phys. Rev. D **71**, 077101 (2005).
- [22] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **76**, 014018 (2007).
- [23] U. Bitenc *et al.* [BELLE Collaboration], Phys. Rev. D **77**, 112003 (2008)
- [24] L. M. Zhang *et al.* [BELLE Collaboration], Phys. Rev. Lett. **96**, 151801 (2006).
- [25] K. Abe *et al.* [BELLE Collaboration], Phys. Rev. Lett. **99**, 131803 (2007).
- [26] P. del Amo Sanchez *et al.* [BABAR Collaboration], Phys. Rev. Lett. **105**, 081803 (2010).
- [27] W. M. Sun (CLEO collaboration), Prepared for 30th International Symposium on Physics in Collision (PIC 2010), Karlsruhe, Germany, 1-4 Sep 2010.
- [28] J. Beringer *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).
- [29] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. Lett. **100**, 061803 (2008).
- [30] M. Staric *et al.* [Belle Collaboration], Phys. Lett. B **670**, 190 (2008).
- [31] T. Aaltonen *et al.* [CDF Collaboration], arXiv:1207.2158 [hep-ex].
- [32] M. Ciuchini, E. Franco, D. Guadagnoli, V. Lubicz, M. Pierini, V. Porretti and L. Silvestrini, Phys. Lett. B **655**, 162 (2007).
- [33] A. L. Kagan and M. D. Sokoloff, Phys. Rev. D **80**, 076008 (2009).

- [34] G. Casarosa [*BABAR* collaboration], Prepared for 5th International Workshop on Charm Physics (Charm 2012), Honolulu, Hawai'i, USA, 14-17 May 2012.
- [35] M. Staric [*Belle* collaboration], Prepared for 5th International Workshop on Charm Physics (Charm 2012), Honolulu, Hawai'i, USA, 14-17 May 2012.